

Volume contents

Volume 48 (1994)

| | | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Issue contents | 1 | M. Meise, W. Janning Localization of thoracic imaginal-disc precursor cells in the early embryo of <i>Drosophila melanogaster</i> | 109 |
| J.B.L. Bard, J.E. McConnell, J.A. Davies Towards a genetic basis for kidney development | 3 | T. Gutjahr, C.E. Vanario-Alonso, L. Pick, M. Noll Multiple regulatory elements direct the complex expression pattern of the <i>Drosophila</i> segmentation gene <i>paired</i> | 119 |
| M. Tio, C. Ma, K. Moses <i>spitz</i> , a <i>Drosophila</i> homolog of transforming growth factor- α , is required in the founding photoreceptor cells of the compound eye facets | 13 | M. Heikinheimo, A. Lawshé, G.M. Shackleford, D.B. Wilson, C.A. MacArthur <i>Fgf-8</i> expression in the post-gastrulation mouse suggests roles in the development of the face, limbs and central nervous system | 129 |
| M. Freeman The <i>spitz</i> gene is required for photoreceptor determination in the <i>Drosophila</i> eye where it interacts with the EGF receptor | 25 | Issue contents | 139 |
| U. Heberlein, A. Penton, S. Falsafi, D. Hackett, G.M. Rubin The C-terminus of the homeodomain is required for functional specificity of the <i>Drosophila</i> <i>rough</i> gene | 35 | J. Song, J.M.W. Slack Spatial and temporal expression of basic fibroblast growth factor (FGF-2) mRNA and protein in early <i>Xenopus</i> development | 141 |
| M. Park, S.A. Monsma, M.F. Wolfner Two tightly-linked <i>Drosophila</i> male accessory gland transcripts with the same developmental expression derive from independent transcription units | 51 | J.C. Ruiz, F.L. Conlon, E.J. Robertson Identification of novel protein kinases expressed in the myocardium of the developing mouse heart | 153 |
| J. Casanova, M. Llimargas, S. Greenwood, G. Struhl An oncogenic form of human <i>raf</i> can specify terminal body pattern in <i>Drosophila</i> | 59 | L. Ho, K. Symes, C. Yordán, L.J. Gudas, M. Mercola Localization of PDGF A and PDGFR α mRNA in <i>Xenopus</i> embryos suggests signalling from neural ectoderm and pharyngeal endoderm to neural crest cells | 165 |
| Issue contents | 65 | A. Brunner, T. Twardzik, S. Schneuwly The <i>Drosophila</i> <i>giant lens</i> gene plays a dual role in eye and optic lobe development: Inhibition of differentiation of ommatidial cells and interference in photoreceptor axon guidance | 175 |
| S. Taraviras, A.P. Monaghan, G. Schütz, G. Kelsey Characterization of the mouse HNF-4 gene and its expression during mouse embryogenesis | 67 | Y.-H. Liu, L. Ma, L.-Y. Wu, W. Luo, R. Kundu, F. Sangiorgi, M.L. Snead, R. Maxson Regulation of the <i>Msx2</i> homeobox gene during mouse embryogenesis: A transgene with 439 bp of 5' flanking sequence is expressed exclusively in the apical ectodermal ridge of the developing limb | 187 |
| J.O. Lewis, S.T. Crews Genetic analysis of the <i>Drosophila</i> <i>single-minded</i> gene reveals a central nervous system influence on muscle development | 81 | | |
| J.A. Helms, S. Kuratani, G.D. Maxwell Cloning and analysis of a new developmentally regulated member of the basic helix-loop-helix family | 93 | | |

| | | | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| G.W. Robinson, K.A. Mahon Differential and overlapping expression domains of <i>Dlx-2</i> and <i>Dlx-3</i> suggest distinct roles for <i>Distal-less</i> homeobox genes in craniofacial development | 199 | G.-Q. Zhao, H. Eberspaecher, M.F. Seldin, B. de Crombrugghe The gene for the homeodomain-containing protein Cart-1 is expressed in cells that have a chondrogenic potential during embryonic development | 245 |
| A. Bartholomä, K.-A. Nave NEX-1: a novel brain-specific helix-loop-helix protein with autoregulation and sustained expression in mature cortical neurons | 217 | B. Christen, M. Bienz Imaginal disc silencers from <i>Ultrabithorax</i> : evidence for <i>Polycomb</i> response elements | 255 |
| Y. Li, M.L. Allende, R. Finkelstein, E.S. Weinberg Expression of two zebrafish <i>orthodenticle</i> -related genes in the embryonic brain | 229 | Author index | 267 |
| | | Subject index | 269 |
| | | Volume contents | 273 |

